

Comment on “Relevant Length Scale of Barchan Dunes”

In a recent experimental breakthrough, Hersen *et al.* [1] demonstrated that by changing the agitating medium from air to water, one can obtain, on laboratory scale, dunes that are downsized copies of desert dunes, thereby overcoming a major obstacle for their systematic study. Here we argue in two steps (i), (ii) that an alternative data analysis leads to some conclusions that are qualitatively and quantitatively different from Hersen *et al.*'s but justify their similarity hypothesis on different grounds.

It has been a long-standing debate whether the various shapes of desert dunes share the scale-invariance of the turbulent wind field that creates them. Careful field measurements of barchan dunes have recently provided compelling evidence that scale invariance is borne out only *transverse* to the wind direction, whereas it is systematically broken *along* the wind direction [2]. That the responsible characteristic scale is the windward distance ℓ_s over which the sand flux adapts to a sudden change in wind speed or sand supply, was already vaguely anticipated by Bagnold. But only recent theoretical developments succeeded in making this notion fully quantitative and established its crucial role for systematic shape variations and shape transitions of aeolian dunes [3].

(i) The analysis of similarities of aeolian and submarine dunes in Ref. [1] suffers from mixing transverse and longitudinal dune dimensions. From what was said above, a genuine test of similarity should address the longitudinal geometry *directly*, thereby avoiding influences [2] from uncontrolled parameters affecting the transverse direction. Recombining the data from Ref. [1], we can infer the (affine) aeolian and submarine height-length (H - L) relations that suggest a considerably larger scale factor than estimated in [1]. Our best estimate for the latter is however obtained from analysing data [1, 4] for the dune migration speed V as follows.

(ii) The (trivial) fact that for a barchan dune, which loses practically no sand over its crest, V is proportional to the flux q_c over the crest divided by its height, is the origin of the popular relation $V \propto H^{-1}$ adopted in Ref. [1]. Note however that the latter would only hold for size-independent q_c , i.e. in the case of scale-invariant dune shape, inconsistent with the observation of broken scale invariance crucially underlying the similarity hypothesis by Hersen *et al.* Yet, one can show on quite general grounds [3] that (essentially) $V \propto L^{-1}$, L being the total length of either smooth heaps or barchan dunes. Fig. 1 demonstrates that this (but *not* $V \propto H^{-1}$) is compatible with *both* aeolian and submarine data. To superimpose the data, lengths had to be rescaled by $6.2 \cdot 10^3$ (in line with our recombined $H - L$ relations) instead of the $1.7 \cdot 10^3$ suggested in [1], and times by $2.3 \cdot 10^6$.

Assuming $\bar{\ell}_s^{\text{air}} \approx 0.8$ m [3] for the flux-weighted average of ℓ_s [5] this implies $\bar{\ell}_s^{\text{water}} \approx 0.13$ mm equal to the grain diameter, as naturally expected for direct fluid entrain-




FIG. 1: Submarine (black [1]) and aeolian (gray [4]; rescaled) migration speed data compared to H^{-1} (dotted [1]) failing for small dunes, and L^{-1} (solid [3]) compatible with the data.

ment. In conclusion, we agree with Ref. [1] that there is an intriguing similarity between submarine and aeolian dunes, but on somewhat different grounds. Further quantitative confirmation of this similarity by data for small submarine dunes, backing the highest submarine data point in Fig. 1, would be desirable.

Klaus Kroy¹ and Xiang Guo²

¹Hahn-Meitner Institut

Glienicker Str. 100, 14109 Berlin, Germany

²Department of Building and Construction, City University of HongKong, HongKong

-
- [1] P. Hersen, S. Douady, and B. Andreotti, Phys. Rev. Lett. **89**, 264301 (2002).
 - [2] G. Sauermann, P. Rognon, A. Poliakov, and H. J. Herrmann, Geomorphology **36**, 47 (2000).
 - [3] G. Sauermann, K. Kroy, and H. J. Herrmann, Phys. Rev. E **64**, 031305 (2001); K. Kroy, G. Sauermann, and H. J. Herrmann, Phys. Rev. Lett. **88**, 054301 (2002), and Phys. Rev. E **66**, 031302 (2002).
 - [4] H. J. Finkel, J. Geol. **67**, 614 (1959).
 - [5] While the average saltation length, the scale l_{drag} introduced in Ref. [1], and $\bar{\ell}_s$ are all proportional to each other on dimensional grounds, their absolute values are generally strongly diverse due to distinctly different dependencies on the wind speed [3], which prohibits their identification even for order-of-magnitude estimates.